

Zooplankton Morphology, Abundance, and Their Relationship with Physicochemical Parameters in Coastal Waters of Fuerte, Caoayan, Ilocos Sur

Jhomar P. Madriaga¹, Antonio N. Ayop²

¹Department of Social Welfare and Development, Provincial Operations Office - Ilocos Sur

²College of Arts and Sciences, University of Northern Philippines, Philippines

¹madriagaj71@gmail.com

²antonio.ayop@unp.edu.ph

ABSTRACT

Zooplankton, which graze on the water column, play a crucial role in maintaining the balance of phytoplankton communities and reducing the frequency of blooms. Despite their substantial contributions to the marine ecosystem, their value to coastal communities was frequently overlooked because of their microscopic nature. The lack of data on marine zooplankton in the Ilocos Sur Region motivated this study conducted in Brgy. Fuerte, Caoayan, Ilocos Sur, to generate new insights into the zooplankton community. It aims to analyze zooplankton morphology, abundance, and physicochemical parameters. Sample collection was conducted during the evening (7 pm to 10 pm). Surface water samples were collected using a handheld fine mesh plankton net (50 µm) and preserved with formaldehyde and Lugol's solution. Thirteen groups of zooplankton populations were recorded in Brgy. Fuerte, Caoayan, Ilocos Sur. They were categorized into two kingdoms: Animalia and Chromista. Common zooplankton included Nauplius larvae, Calanus sp. (Calanoid), and Dioithona sp. (Cyclopoid). The overall density of zooplankton recorded was 192,050 cells/L, which comprises four stations. Regarding species density, the top five dominant species were Favella sp. (Tintinnids) with 163,689 cells/L, Nauplius larvae with 13,478 cells/L, Calanus sp. (Calanoid) with 9,300 cells/L, Sabellaria sp. (Middle trochophore larvae) with 1,283 cells/L, and Dioithona sp. (Cyclopoid) with 1,211 cells/. Future studies should investigate the complex relationships between physicochemical factors and zooplankton abundance and diversity while extending the sampling period to include additional zooplankton species. Understanding these relationships will enhance the comprehension of ecosystem dynamics and develop more effective coastal water management strategies.

Keywords: Zooplankton, Stations, Density, Species diversity, Water parameters, Population, Ilocos Sur

INTRODUCTION

Marine ecosystems support a wide variety of animals, from microscopic plankton to enormous whales, and encompass a variety of habitats like coral reefs, coastal regions, and open oceans (Ilac et al., 2024; Mendoza et al., 2023). Central to these ecosystems are zooplankton, which play an integral role in nutrient cycling and act as a foundational food

source for many marine organisms, including fish and whales. They play a vital function in preserving the equilibrium and general health of the marine environment.

Spanning a wide size range—from microscopic protozoans to larger metazoans such as jellyfish—zooplankton are indispensable contributors to the food web and nutrient cycling, underscoring their ecological importance. They contribute to ecological processes such as carbon sequestration through the biological carbon pump, a mechanism that aids in transporting carbon to deeper ocean layers (Turner, 2004). Their role in regulating phytoplankton populations also helps maintain water quality and balance within aquatic ecosystems. Notably, they facilitate carbon sequestration through the biological carbon pump, as elucidated by Pinti et al. (2023), and enhance carbon export through fecal pellet deposition (Halfter et al., 2020). Furthermore, their ability to regulate phytoplankton populations contributes to preserving water quality and ecological balance (Okogwu, 2010; Turner, 2004).

The abundance, diversity, and biomass of zooplankton profoundly influence aquatic environments. As biomonitoring indicators, they serve as reliable indicators of ecosystem health (Davies et al., 2009; Florendo, 2003). Copepods, a subclass of zooplankton, exhibit filter-feeding capabilities and contribute to water quality maintenance. However, they are susceptible to ecological disturbances, such as pollution, which can disrupt metabolic processes (Lauritano et al., 2012). In the context of fisheries, zooplankton provide a nutrient-rich and cost-effective protein source for fish, thereby supporting global economies (El-Fattah et al., 2008; Kibria et al., 1997). For instance, (Mamaril, 2001), in Lake Taal in Batangas, renowned for its highly valued fish "*Sardinella tawilis*," zooplankton plays a crucial role as food items during early life and some adult stages.

Environmental factors like temperature, salinity, and human activity influence the distribution of zooplankton worldwide. With the help of tropical waters and nutrient-rich upwellings, zooplankton flourish in ecosystems like coral reefs and mangroves in the Philippines, known for its rich marine biodiversity (Villanoy et al., 2011). Nevertheless, seasonal variations and anthropogenic disturbances often lead to fluctuations in local populations (David et al., 2005). Advanced research techniques, including water quality monitoring system tools (Divina et al., 2023), in-situ sampling, and satellite remote sensing offer important insights into zooplankton dynamics (Basedow et al., 2019). Zooplankton form and distribution are influenced by biological succession stages, waterbody size, and trophic circumstances (Cloern, 2001). These studies improve resource management techniques and advance our understanding of their ecological significance (Hays et al., 2005; Villanoy et al., 2011). Environmental pressures endangering zooplankton populations globally include pollution, overfishing, and climate change. Their ability to survive and procreate is hampered by ocean acidification, warming temperatures, and microplastic pollution, which upsets marine food webs (Cole et al., 2013; Fabry et al., 2008). Global collaboration is required to

The Vector: International Journal of Emerging Science, Technology and Management
Volume 33, Issue 1, January - December 2024

monitor and lessen human impacts on ocean health and address these issues (Richardson, 2008). Pollution and climate change in the Philippines exacerbate the problems that marine ecosystems confront, causing zooplankton populations to drop and endangering the ecosystem's health (Botterell et al., 2023; Lasco, 2022; Shaira Elyza R. et al., 2024).

The Ilocos Sur Region lacks published data on marine zooplankton studies, making this potentially the area's first documented account of zooplankton communities. The coastal waters of Brgy. Fuerte, Caoayan, Ilocos Sur, are an ideal site for zooplankton research due to their rich fishing grounds and distinctive features, including an oyster farm, estuarine regions, Fuerte Beach (a residential area), and Choco Surf (a tourist destination). These diverse environments and varying levels of human and natural disturbances provide a valuable opportunity to examine their effects on zooplankton populations.

Objectives of the Study

This study examined the morphological characteristics, abundance, diversity, and the relationship between physicochemical parameters across four selected stations in Brgy. Fuerte, Caoayan, Ilocos Sur.

METHODOLOGY

Research Design

This study used a mixed design (qualitative-descriptive) to determine zooplankton species composition, abundance, and diversity at four stations in Barangay Fuerte, Caoayan, Ilocos Sur.

Study Site

The study was conducted in Barangay Fuerte, Caoayan, Ilocos Sur. Fuerte is a barangay in the municipality of Caoayan, located in the province of Ilocos Sur.

The 2020 Census (PhilAtlas, 2024) reported a population of 2,675, representing 13.67% of Caoayan's total population. Four coastal stations were selected: ST1-Choco Surf (tourist area), ST2-Fuerte Beach (residential area), ST3-Estuary, and ST4-Oyster Farm. Each station has unique characteristics, including anthropogenic disturbances that impact the zooplankton community. The coastal waters of Fuerte Caoayan are also fishing grounds for marine shore fish and shellfish.

Data Gathering Procedure

Sample Collection. Sampling was conducted from 7:00 PM to 10:00 PM to account for the nocturnal behavior of zooplankton, which undergoes vertical migration at night. From February to March 2024, which serves as cool-dry to warm-dry months, samples were

collected three times over a month, with a 10-day interval between each collection. Surface samples were collected using a handheld fine mesh plankton net (50 μm) and a 10L bucket. Water was filtered through the net, washed, and transferred into 100ml empty canisters. Samples were labeled with the date and time of sampling. Three canisters were collected from each station, representing three replicates.



Figure 1
Map showing the study site. A) Map of the Philippines showing Region 1. B) Map of Region 1 showing Brgy. Fuerte, Caoayan. C) Map of Brgy. Fuerte, Caoayan showing the four sampling stations (red circles).

Zooplankton Samples Preservation. Collected samples were preserved with 5% formaldehyde for 1 minute. Lugol's solution was added to stain cells using a medicine dropper and then mixed gently. Phytoplankton samples were undistributed in the dark at room temperature for 48 hours to settle. After sedimentation, the water bottle was carefully sucked out, and the final volume was adjusted to 50 mL. The remaining 50mL water sample was used for cell count and zooplankton identification. After the experiment, the water samples were disposed of properly in the designated waste containers. Used bottles, containers, and other materials were also discarded properly.

Taxonomic Identification. The volume of collected samples from each station was standardized to 50mL. Gently shake the canister and mix the upper and lower layers. Five mL aliquots were examined under a microscope using a 1ml Sedgewick-R after the counting chamber. Zooplankton identification at the genus level and phylum categorization was based on morphology, structures, and shape, aided by manuals and field guides of Slotwinski et al. (2014); Yamani et al. (2011a) & (2011b). Photographs were taken at 4x and 10x magnification.

Zooplankton Density. Aliquots were placed in the Sedgewick-Rafter cell under the microscope for accurate phytoplankton counting. To prevent overflowing, coverslips were used to mitigate evaporation-induced air space formation. The counting process involved moving the cell vertically along the first column of squares, tallying organisms in each square. The total plankton count in a 50 ml water sample was determined using the formula established by (Santhanam et al., 1989).

$$N=(n \times v)/V \quad (1)$$

Where in: N = the total number of plankton cells per liter of water-filtered; n = average number of plankton cells in 1 ml. of plankton samples; v = volume of plankton concentration (ml); V = volume of water filtered (l)

Species Diversity. Zooplankton species diversity was determined using the Shannon-Weiner Index, which is presented below;

$$H' = - \sum_{n-1}^n (p_i * \ln p_i) \quad (2)$$

Where pi = proportion of total sample represented by species; n = represents the total number of species (or different categories) in the community or sample.

Furthermore, the results of the diversity index were interpreted using the guidelines of Fernando et al. (1998), as cited by (Coracero et al., 2020).

Table 1
Categories of Species Diversity Index

Relative Values	H' Values	Evenness Index
Very High	3.5 and above	0.75 – 1.00
High	3.0 – 3.49	0.5 - 0.74
Moderate	2.5 – 2.99	0.25 - 0.49
Low	2.0 – 2.49	0.15 - 0.24
Very Low	1.9 and below	0.05 - 0.14

Dana Analysis

Mean was used to determine the average value of water parameters and the abundance of zooplankton species.

RESULTS AND DISCUSSIONS

1. Morphological Characteristics of Zooplankton

Thirteen species of zooplankton populations were recorded in Brgy. Fuerte, Caoayan, Ilocos Sur. They were categorized into two kingdoms: Animalia and Chromista. The populations were divided into five phyla with corresponding classes: Arthropoda (three classes: Malacostraca, Copepoda, Branchiopoda), Annelida (one class: Polychaeta), Cnidaria (two classes: Scyphozoa, Cubozoa), Ciliophora (one class: Oligotrichea), and Foraminifera (one class: Globothalamea); see below the taxonomic classification system.

Kingdom: Animalia

Phylum: Arthropoda (Gravenhorst, 1843)

Subphylum: Crustacea (Brünnich, 1772)

Nauplius larvae (**Figure 2a**)

Class: Malacostraca (Latreille, 1802)

Order: Amphipoda (Latreille, 1816)

Family: Gammaridae (Latreille, 1802)

Genus: *Gammarus* (Fabricius, 1775)

Gammarus sp. (**Figure 2b**)

Order: Cumacea (Krøyer, 1846)

Family: Leuconidae (Sars, 1878)

Genus: *Nannastacus* (Bate, 1865)

Nannastacus sp. (**Figure 2c**)

Class: Copepoda (Edwards, 1840)

Order: Calanoida (Sars, 1903)

Family: Calanidae (Dana, 1849)

Genus: *Calanus* (Leach, 1816)

Calanus sp. (**Figure 2d**)

Order: Cyclopoida (Burmeister, 1834)

Family: Oithonidae (Dana, 1853-1855)

Genus: *Dioithona* (Kiefer, 1935)

Dioithona sp. (**Figure 2e**)

Order: Harpacticoida (Sars, 1903)

Family: Ectinosomatidae (Sars, 1903)

Genus: *Microsetella* (Brady & Robertson, 1873)

Microsetella sp. (**Figure 2f**)

Class: Branchiopoda (Latreille, 1817)

Order: Onychopoda (Sars, 1865)

Family: Podonidae (Mordukhai-Boltovskoi, 1968)

Genus: *Evadne* (Lovén, 1836)

Evadne sp. (**Figure 2g**)

Phylum: Annelida (Lamarck, 1802)

Class: Polychaeta (Grube, 1850)

Order: Canalipalpata

Family: Sabellariidae (Johnston, 1865)

Genus: *Sabellaria* (Lamarck, 1818)

Sabellaria sp. (Early trochophore) (**Figure 2h**)

Sabellaria sp. (Mid trochophore) (**Figure 2i**)

Phylum: Cnidaria (Hatschek, 1888)

Class: Scyphozoa (Goette, 1887)

Ephyra (Scyphozoa) (**Figure 2j**)

Class: Cubozoa (Werner, 1973)

Medusa (Cubozoa) (**Figure 2k**)

Kingdom: Chromista

Phylum: Ciliophora (WoRMS, 2024)

Class: Oligotrichea (Bütschli, 1887)

Order: Choreotrichida (Small & Lynn, 1985)

Family: Ptychocylididae (Kofoid & Campbell, 1929)

Genus: *Favella* (Jørgensen, 1924)

Favella sp. (**Figure 2l**)

Phylum: Foraminifera (d'Orbigny, 1826)
Class: Globothalamea (Pawlowski et al., 2013)
Order: Rotaliida (Lankester, 1885)
Family: Discorbidae (Ehrenberg, 1838)
Genus: *Discorbis* (Lamarck, 1804)
Discorbis sp. (**Figure 2m**)

The nauplius larvae, shown in Figure 2a, were found at all stations, with a mean length of $26.22 \mu\text{m} \pm 3.61 \mu\text{m}$. They have three pairs of appendages: the first antennae, the second antennae, and the mandibles. The body is unsegmented, and they have a single eye in the center of the head. (Martin & Davis, 2001).

The amphipod genus *Gammarus* in Figure 2b was found only in Oyster Farm, averaging $82.29 \mu\text{m} \pm 5.46 \mu\text{m}$ in length. They can infiltrate and colonize environments due to their extensive trophic repertoire, foraging adaptability, migration, and tendency to wander. *Gammarus* also have a high reproductive capacity, producing many broods, offspring, and a relatively long lifespan of 1-2 years (Gerhardt et al., 2011).

The genus *Nannastacus* in Figure 2c, found only in Residential, has an average length of $38.86 \mu\text{m} \pm 1.20 \mu\text{m}$. It has elongated bodies segmented into distinct parts, including a carapace fused to the first thoracic segment, forming a protective hood. Cumaceans usually lie submerged in the sediment, needing to stay in contact with the water to pump oxygenated water over their gills (Schram & Koenemann, 2022b).

The calanoid genus *Calanus* in Figure 2d was found at all stations, with a mean length of $88.04 \mu\text{m} \pm 14.22 \mu\text{m}$. They have elongated bodies and segmented appendages for swimming and feeding (Blaxter et al., 1998).

The Cyclopoid genus *Dioithona* in Figure 2e was found at all stations Choco Surf, Residential, Estuary, and Oyster Farm, averaging $103.11 \mu\text{m} \pm 26.28 \mu\text{m}$. They have streamlined bodies, single eyes, antennae, and swimming legs for rapid movement, often as freshwater and marine predators (Schram & Koenemann, 2022a).

Microsetella, a harpacticoid genus, shown in Figure 2f was found at three stations: Fuerte Beach, Estuary, and Oyster Farm with a mean length of $63.56 \mu\text{m} \pm 15.46 \mu\text{m}$. It has an elongated worm-like body equipped with caudal setae that help slow down its sinking velocity (Uye et al., 2002).




Figure 2

(a) Nauplius larvae, (b) *Gammarus* sp., (c) *Nannastacus* sp., (d) *Calanus* sp., (e) *Dioithona* sp., (f) *Microsetella* sp., (g) *Evadne* sp., (h) *Sabellaria* sp. (Early trochophore), (i) *Sabellaria* sp. (Mid trochophore), (j) *Ephyra* (Scyphozoa), (k) *Medusa* (Cubuzoa), (l) *Favella* sp., (m) *Discorbis* sp.

The cladoceran genus *Evadne* in Figure 2g was found only in Estuary, averaging $17.80 \mu\text{m} \pm 0.06 \mu\text{m}$ in length. *Evadne* are oval-shaped with large compound eyes, several swimming appendages, and a long tailspin (Aquascope, 2000).

The tubeworm genus *Sabellaria* (early trochophore) larvae in Figure 2h were found in Estuary and Oyster Farm, averaging $15.62 \mu\text{m} \pm 0.59 \mu\text{m}$. The turf, which beats slowly from one side of the episphere, is wrinkled in the apical and hypospherical regions of the vitelline membrane (Smith & Chia, 2011).

Mid trochophore larvae, shown in Figure 2i, found in Choco Surf, Fuerte Beach, and Oyster Farm average $24.32 \mu\text{m} \pm 2.09 \mu\text{m}$ in length. It is distinguished by a pygidium, a provisional setae, and the hyposphere's division into two segments. The apical turf comprises three bundles of short, laterally beating cilia (Smith & Chia, 2011).

The Ephyra (scyphozoa) in Figure 2j, a jellyfish genus, was found at three stations: Fuerte Beach, Estuary, and Oyster Farm. The mean length of *Ephyrae* was $20.40 \mu\text{m} \pm 0.70 \mu\text{m}$. They have bell-shaped bodies with trailing tentacles, arising from the medusa stage. *Ephyrae* lack the polyp stage, possessing simple radial symmetry and a gelatinous mesoglea between two epithelial layers (Gershwin, 2016).

The Medusa (cubozoan) in Figure 2k, found only in the Estuary, has a mean length of $43.61 \mu\text{m} \pm 0.26 \mu\text{m}$. Box jellyfish, as they're known, have a distinctive cuboidal bell with four tentacles at each corner and a centrally located cluster of appendages called rhopalia containing eyes and sensory structures. This distinguishes them from other jellyfish and reflects their advanced predatory capabilities in the ocean (Gershwin, 2016).

The *Favella* genus in Figure 2l, found in Estuary and Oyster Farms, has a mean length of $16.19 \mu\text{m} \pm 0.93 \mu\text{m}$. It's a hyaline, particle-free lorica that differs in shape from tintinnids. The *Favella* form is predominantly cylindrical with a bowl-shaped body and an aboral constriction (Kim et al., 2010).

Discorbis, a foraminifera genus, shown in Figure 2m was found only on Oyster Farms. It has a mean length of $10.10 \mu\text{m} \pm 0.09 \mu\text{m}$. *Discorbis* are known for their spiral-shaped, chambered shells, commonly found in marine sedimentary environments (Loeblich & Tappan, 1988).

All organisms in Kingdom Animalia that are planktonic during their lives are part of zooplankton communities. Some settle as meroplankton during adulthood. In Kingdom Chromista, organisms such as holoplankton are planktonic for life. They lack chloroplasts or are heterotrophic.

2. Abundance and Diversity of Zooplankton

Table 2 shows the overall zooplankton density in Brgy. Fuerte, Caoayan, Ilocos Sur, comprising four stations. The highest density was in Estuary (159,517 cells/L), followed by Oyster Farm (24,583 cells/L) and Choco Surf (3,311 cells/L). *Favella* sp. (Tintinnids) dominated

The Vector: International Journal of Emerging Science, Technology and Management
Volume 33, Issue 1, January - December 2024

with 163,689 cells/L, followed by Nauplius larvae (13,478 cells/L), *Calanus* sp. (Calanoid) (9,300 cells/L), *Sabellaria* sp. (Middle trochophore larvae) (1,283 cells/L), and *Dioithona* sp. (Cyclopoid) (1,211 cells/L).

Table 2

Zooplankton Species Abundance in the Four Stations of Fuerte, Caoayan, Ilocos Sur

Zooplankton Population	Coastal Waters				Overall Density
	Choco Surf	Fuerte Beach	Estuary	Oyster Farm	
	\bar{x} Density (cells/L)				Σ Density (cells/L)
I. Kingdom Animalia					
1. Nauplius larvae	633	2,144	7,866	2,833	13,478
2. <i>Gammarus</i> sp.				333	333
3. <i>Nannastacus</i> sp.		100			100
4. <i>Calanus</i> sp.	2,133	1,211	2,422	3,533	9,300
5. <i>Dioithona</i> sp.	244	300	466	200	1,211
6. <i>Microsetella</i> sp.		183	133	233	550
7. <i>Evadne</i> sp.			133		133
8. <i>Sabellaria</i> sp. (Early trochophore larvae)			350	656	1,006
9. <i>Sabellaria</i> sp. (Mid trochophore larvae)	300	267		717	1,283
10. Ephyra (scyphozoa)		100	167	133	400
11. Medusa (cubozoan)			100		100
12. <i>Favella</i> sp.			147,544	16,144	163,689
13. <i>Discorbis</i> sp.			333	133	467
Grand Total	3,311	4,306	159,517	24,583	192,050
Shannon's index (H')	0.98	1.28	0.32	1.03	0.63
Shannon evenness (E_{sh})	0.77	0.80	0.18	0.56	0.25
H' Condition Index	Very Low	Very Low	Very Low	Very Low	Very Low

Favella sp. dominated the abundance, with 147,544 cells/L in the Estuary and 16,144 cells/L in the Oyster Farm. Temperature and the summer period contributed to its high abundance at both stations. Similar findings were observed in a study by Durmus et al. (2023), where Tintinnids were prevalent throughout spring and summer, with temperature as the primary factor influencing their composition.

Nauplius larvae, commonly found in all stations, had the highest density in the Estuary (7,866 cells/L), followed by Oyster Farm (2,833 cells/L), and Choco Surf (633 cells/L). This larval stage is characteristic of most crustaceans. (Naung, 2018) noted copepod and barnacle nauplii in shallow coastal estuaries. *Calanus* sp. had the highest density in Oyster Farms (3,533 cells/L), followed by the Estuary (2,422 cells/L), and Fuerte Beach (1,211 cells/L). *Calanus*, a vital ecosystem component and planktivorous fish food mentioned by (Ramírez & Sabatini, 2000), is also temperature-dependent, especially in summer.

Sabellaria sp. (middle trochophore) larvae were the second most abundant species at all three stations, except the Estuary. Oyster Farm had the highest density (717 cells/L), followed by Choco Surf (300 cells/L), and Fuerte Beach (267 cells/L). These larvae are just before settlement. The Estuary's narrow inlet and restricted water flow during low tide made it unsuitable for settlement. The other stations provided favorable conditions due to their wide areas and suitable substrates. Larvae exhibit selective settling in areas with appropriate sediment conditions (Tait & Dipper, 1998). *Dioithona* sp. was found at all stations, with the highest density at the Estuary (466 cells/L), followed by Fuerte Beach (300 cells/L), and the lowest at Oyster Farm (200 cells/L). This zooplankton group belongs to the Orders Cyclopoida and Calanoida, indicating suitable marine water conditions for survival (Canencia, 2017).

The Estuary station has the highest overall zooplankton density, with a mean density of 159,517 cells/L. The Oyster Farm station has the second highest density, at 192,050 cells/L. The Fuerte Beach station has the third highest density, at 4,306 cells/L. The Choco Surf station has the lowest density, at 3,311 cells/L. The high density in the Estuary is due to the nutrient-rich environment, which fosters optimal plankton growth. Estuaries receive nutrients from land and ocean sources, promoting plankton proliferation (Cloern, 2001). Mixing fresh and saltwater creates a dynamic ecosystem supporting diverse plankton populations. The elevated plankton levels in the Estuary align with Cloern's understanding of estuarine ecology.

In contrast, the other stations are near residential zones and tourist areas that could affect zooplankton populations. Human activities and pollution inputs, such as garbage, can cause a decline in marine zooplankton abundance (Thushari & Senevirathna, 2020). With their high human activity, Choco Surf and Fuerte Beach experienced lower zooplankton populations due to these factors.

Pollution in the Bilbao estuary enhanced zooplankton abundance in the outer euhaline zone but limited it in the inner zone, as observed by (Uriarte & Villate, 2004). (Mwagana et al., 2018) further demonstrated that water pollution, particularly from total

nitrogen and nitrate, significantly influenced zooplankton biodiversity, though the study did not test this. (Echeveste et al., 2011) highlighted the toxic effect of complex organic pollutants on phytoplankton, a crucial food source for zooplankton. These studies collectively underscore the detrimental impact of human activities and pollution on marine zooplankton abundance.

The overall species diversity of Zooplankton populations is very low, with $H' < 1.99$ (Table 1 & 2). Evenness (E_{Sh}) is slightly even at Fuerte Beach ($E_{Sh} = 0.80$) and Choco Surf ($E_{Sh} = 0.77$), but not at Oyster Farm ($E_{Sh} = 0.25$), and Estuary ($E_{Sh} = 0.56$) (Table 1 & 3). Shannon's theorem states that evenness (E_{Sh}) ranges from 0 to 1, with 1 representing high evenness.

3. Significant Difference Between and Among the Abundance and Species of Zooplankton in the Different Sampling Stations

Table 3 shows the ANOVA results for zooplankton abundance across different stations. The stations revealed no substantial variation in the abundance of the species across different stations because the upper and lower limits of all species overlapped in the 95% confidence interval. Variations across species demonstrated significant differences based on Tukey's Honestly Significant Difference. In the group of *Favella* sp.-Calanoid ($p = 0.000143$), *Favella* sp.-Cyclopoid ($p = 0.0000349$), *Favella* sp.-Early trochopore larvae ($p = 0.00141$), *Favella* sp.-Ephyra scyphozoa ($p = 0.0135$), Harpacticoid-*Favella* sp. ($p = 0.00131$), Mid trochophore larvae-*Favella* sp. ($p = 0.00399$), and Nauplius larvae-*Favella* sp. ($p = 0.0000915$).

Table 3

One-way Analysis of Variance (One-Way ANOVA) on the significant difference in zooplankton abundance

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
Species	12	1.056e+11	8.802e+09	3.129	0.000494	***
Station	3	1.454e+10	4.848e+09	1.723	0.164175	
Residuals	167	4.698e+11	2.813e+09			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

The high abundance of *Favella* sp. was due to their exhibits of behavioral selective particle feeding mechanisms, which are common to the group of tintinnid ciliate, according to (Stoecker et al., 1995). It also exhibits intricate behavioral responses to phytoplankton, such

as adjustments in swimming speed and turning rate when interacting with its exudates (Buskey & Stoecker, 1989). On the other hand, harpacticoid copepods, along with their nauplii, are essential components of marine ecosystems. They possess varied life cycles, progressing through six nauplius and six copepodid stages, and are well-suited to various habitats, spanning from marine to freshwater environments (Dahms & Qian, 2004). Lastly, the abundance of this mid-trochophore larva is due to its dominant larval stages in different phyla, which constitute the major branch of Bilateria. (Dahms & Qian, 2004).

Coastal Water Parameters

Table 4 shows the mean values of water parameters at the four stations in Fuerte, Caoayan, and Ilocos Sur compared with the standard values of seawater and estuarine water.

Coastal water salinity varies widely, with Choco Surf recording the highest average of 32.77‰ (± 0.03), followed by Fuerte Beach at 32.40‰ (± 0.60). The Estuary has a lower average of 29.67‰ (± 0.58), while the Oyster Farm has the lowest at 28.63‰ (± 0.58). All stations met the optimum marine biota salinity range of 0.5 ‰-40 ‰ (Jain, 2011; van Velzen, 2022). Choco Surf and Fuerte Beach are euhaline (32.77 ± 0.03 ‰), while Oyster Farm and Estuary are polyhaline (18 ‰-30 ‰) based on Mitsch and Gosselink (1986) as cited by Ohrel and Register (2006). Mixing freshwater from the Mestizo River slightly decreased salinity in these stations. Salinity, a chemical stressor of dissolved salts, indicates species tolerance or range limitations and affects community organization (Smyth & Elliott, 2016).

Table 4 shows that all stations maintain optimal temperatures between 25-31°C (DENR Administrative Order 2016-08, 2016). Oyster Farm and Estuary recorded slightly higher temperatures (28.23 ± 0.28 and 28.10 ± 0.40 °C, respectively) than Fuerte Beach (27.30 ± 0.15 °C) and Choco Surf (27.27 ± 0.13 °C). The colder sea surface temperature and winds during nighttime sampling may have affected water cooling.

Coastal waters exhibit varying pH levels, with the highest average pH (8.14 ± 0.08) in the Estuary, which is slightly alkaline. Fuerte Beach and Choco Surf have similar pH levels (7.93 ± 0.03 and 7.90 ± 0.03 , respectively). The Oyster Farm area has the lowest average pH (7.94 ± 0.06), slightly acidic. All stations' pH levels are optimal (6.50-8.50) for marine and estuarine water. The estuary's slightly higher pH (8.14 ± 0.08) is due to the mixing of freshwater and saltwater at the mouth, as suggested by the (United States Environmental Protection Agency, 2006). This could include the estuarine mouth, which connects to the ocean.

Across all stations, turbidity was lower than the optimum requirements in Marine and Estuarine (Table 4). Sandy substrates at Choco Surf and Fuerte Beach settle sediment-granulated particles quickly, even under strong wave action. In the Estuary and Oyster Farm, sandy-muddy substrates with less water movement and agitation exhibit low sedimentation. Lower turbidity reduces pollutant concentration, promoting a healthy ecosystem (Fondriest Environmental Inc., 2014).

Table 4
Mean Physicochemical Parameters in the four stations

Water Parameters	Coastal Waters				Optimum Requirements	
	Choco Surf	Fuerte Beach	Estuary	Oyster Farm	Marine	Estuarine
	\bar{x} values					
a. Salinity (‰)	32.77±0.03	32.40±0.60	29.67±0.58	28.63±0.58	30-40 (Jain, 2011)	0.5-30 (van Velzen, 2022)
b. Temperature (°C)	27.27±0.13	27.30±0.15	28.10±0.40	28.23±0.28	25.00-31.00 (DAO, 2016)	0-30 (Kennish, 1990)
c. pH	7.90±0.03	7.93±0.03	8.14±0.08	7.94±0.06	6.50-8.50 (DAO, 2016)	7.0-7.5 (USEPA, 2006)
d. Turbidity (NTU)	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	30 (FEI, 2014))	1-100 (Gallegos et al. 1990)
e. Dissolved oxygen (mg/L)	8.90±1.35	8.93±1.33	4.97±0.26	5.50±1.04	> 5.00 (DAO, 2016)	5.5-11.8 (Priya et al., 2022)
f. Total Dissolved Solids (ppm)	266.33±0.67	267.33±1.20	241.00±5.03	234.30±5.33	>10,000 (Adjovu et al., 2023)	500-30,000 (Horsburgh & Wilson, 2007)
g. Electrical conductivity (mS/cm)	53.43±0.07	53.73±0.26	48.63±0.88	47.40±1.02	30.00-60.00 (Zheng et al., 2018)	35.10-49.34 (Cameron & Pritchard, 1963)

Coastal waters vary in dissolved oxygen levels. Choco Surf and Fuerte Beach have the highest average concentrations (8.90±1.35 mg/L and 8.93±1.33 mg/L, respectively), while the Estuary and Oyster Farm areas have lower concentrations (4.97±0.26 mg/L and 5.50±1.04 mg/L, respectively). These levels meet the optimum requirement except in the Estuary. Choco Surf and Fuerte Beach exceed the requirement of > 5 mg/L, while the Estuary falls below it. This oxygen deficiency could be due to rapid oxygen consumption, especially at night, caused by respiration and organic matter degradation (Horak et al., 2016).

Coastal waters exhibit significant TDS variation, with Fuerte Beach having the highest average concentration (267.33±1.20 ppm) and Choco Surf (266.33±0.67 ppm) closely behind. The Estuary (241.00±5.03 ppm) shows a decrease, while the Oyster Farm area (234.30±5.33 ppm) has the lowest. All stations below the optimum TDS requirements (500 to > 10,000 ppm) (Adjovu et al., 2023; Horsburgh & Wilson, 2007). Lower TDS indicates minimal nutrient influx, resulting in fewer ions and organic ions. Weber-Scannell and Duffy (2007) suggest alterations in ionic composition can eliminate certain species while promoting others, but it doesn't

quantify TDS and can't determine if species composition has been compromised. The area's good condition is attributed to fewer ions and no signs of eutrophication.

Coastal waters exhibit variations in electrical conductivity. Fuerte Beach and Choco Surf have the highest average conductivity (53.73 ± 0.26 mS/cm and 53.43 ± 0.07 mS/cm, respectively). In contrast, the Estuary and Oyster Farm areas have lower values (48.63 ± 0.88 mS/cm and 47.40 ± 1.02 mS/cm, respectively). Based on (Chanson et al., 2005; Zheng et al., 2018), all values fall within the optimum range of 30.00-60.00 or 35.10-49.34 mS/cm. According to (Aluwong et al., 2024), increased electrical conductivity indicates elevated dissolved solids, which can harm aquatic organisms, human well-being, and industrial operations. However, the EC levels in this study were not alarming as they met the optimum requirement, suggesting low inorganic and organic ions. Despite the low dissolved solids, the EC remained within the optimum range due to the concentration of dissolved salts.

CONCLUSIONS

The findings of the study revealed that Brgy. Fuerte, Caoayan, Ilocos Sur exhibited a high species richness with a distinct group of morphological characteristics of a planktonic microbiota. Choco Surf Point exhibited the highest salinity, temperature, dissolved solids, and electrical conductivity, while Fuerte Beach recorded the highest pH and dissolved oxygen levels among the studied stations. The highest recorded physicochemical values of the two stations are attributed to frequent water movement, such as wave action. However, Brgy. Fuerte, Caoayan, Ilocos Sur, and the four sampling stations are in very low conditions for species diversity, with no evenness in a population due to the significant dominance of the species *Favella* sp.

RECOMMENDATIONS

Implementing a comprehensive monitoring program to evaluate nutrient water quality in Brgy. Fuerte, Caoayan, and Ilocos Sur is imperative to ensure consistent adherence to minimum standards and identify potential alterations or trends over time. The findings of this study will be disseminated to the community for ongoing public awareness. Investigate the diel vertical migration pattern of zooplankton to elucidate the movement of zooplankton to deeper depths as they ascend from the surface.

ETHICAL STATEMENT

The study underwent Harmonized Gender and Development Guidelines (HGDG) and the College of Arts and Sciences Research Ethics Committee (CAS RERC certification no. 2024-

The Vector: International Journal of Emerging Science, Technology and Management
Volume 33, Issue 1, January - December 2024

015). The researcher asked permission from the local government unit (LGU) of Fuerte, Caoayan, Ilocos Sur, to collect water samples in study sites through a requested letter. Also, the study was at low risk to the researcher's health since it had minimal fixative use for water samples. After being used, the water samples were properly disposed of following the laboratory protocol.

ACKNOWLEDGMENT

This undertaking is impossible without the support of individuals, particularly the Government Unit of Caoayan, Ilocos Sur, headed by Hon. Mayor Germelina Singson Goulart, who granted permission to conduct the investigation within their coastal water. We are grateful to our research panelists, Dr. Adora G. Ilac, Mrs. Joanna Mae S. Foronda, and Prof. Jessy C. Domingo, for their valuable insights and expertise on the study. We also thank our statistician, Ms. Roselyn Rabanzo, for providing statistical support and expertise. Additionally, we acknowledge the assistance of the Laboratory Science Office, headed by Prof. Jonnel Torres, for providing the necessary laboratory facilities and resources to complete the study.

REFERENCES

- Adjovu, G. E., Ali Shaikh, T., Stephen, H., & Ahmad, S. (2023). Utilization of machine learning models and satellite data for the estimation of total dissolved solids in the Colorado River system. *World Environmental and Water Resources Congress 2023: Adaptive Planning and Design in an Age of Risk and Uncertainty*, 1147–1160. <https://doi.org/10.1061/9780784484852.105>
- Aluwong, K. C., Hashim, M. H. M., Ismail, S., & Shehu, S. A. (2024). Modeling pH changes and electrical conductivity in surface water as a result of mining activities. *Naukovyi Visnyk Natsionalnoho Hirnychoho Universytetu*, 1, 122–131. <https://doi.org/10.33271/nvngu/2024-1/122>
- Aquascope. (2000). Cladocerans of the *Evadne* family. *Tjärnö Marine Biological Laboratory, Strömstad, Sweden*. <https://www.vattenkikaren.gu.se/fakta/arter/crustace/diplostr/evadne/evade.html>
- Basedow, S. L., McKee, D., Lefering, I., Gislason, A., Daase, M., Trudnowska, E., Egeland, E. S., Choquet, M., & Falk-Petersen, S. (2019). Remote sensing of zooplankton swarms. *Scientific Reports*, 9(1), 1–10. <https://doi.org/10.1038/s41598-018-37129-x>
- Blaxter, J. H. S., Douglas, B., Tyler, P. A., & Mauchline, J. (1998). The biology of calanoid copepods. *Advances in Marine Biology*, 33, 1–710. [https://doi.org/10.1016/S0065-2881\(08\)60212-9](https://doi.org/10.1016/S0065-2881(08)60212-9)

- Botterell, Z. L. R., Lindeque, P. K., Thompson, R. C., & Beaumont, N. J. (2023). An assessment of the ecosystem services of marine zooplankton and the key threats to their provision. *Ecosystem Services*, *63*, 101542. <https://doi.org/10.1016/j.ecoser.2023.101542>
- Buskey, E. J., & Stoecker, D. K. (1989). Behavioral responses of the marine tintinnid *Favella* sp. to phytoplankton: Influence of chemical, mechanical and photic stimuli. *Journal of Experimental Marine Biology and Ecology*, *132*(1), 1–16. [https://doi.org/10.1016/0022-0981\(89\)90173-1](https://doi.org/10.1016/0022-0981(89)90173-1)
- Canencia, M. O. P. (2017). Marine zooplankton distribution model and seriation index across different habitat types. *International Journal for Research in Applied Science and Engineering Technology*, *5*(11), 1646–1692. <https://doi.org/10.22214/ijraset.2017.11236>
- Chanson, H., Brown, R., Ferris, J., Ramsay, I., & Warburton, K. (2005). Preliminary measurements of turbulence and environmental parameters in a sub-tropical estuary of Eastern Australia. *Environmental Fluid Mechanics*, *5*(6), 553–575. <https://doi.org/10.1007/s10652-005-0928-y>
- Cloern, J. E. (2001). Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology Progress Series*, *210*, 223–253. <https://doi.org/10.3354/meps210223>
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., & Galloway, T. S. (2013). Microplastic ingestion by zooplankton. *Environmental Science & Technology*, *47*(12), 6646–6655. <https://doi.org/10.1021/es400663f>
- Coracero, E. E., Jr., & Mariano, P. L. M. (2020). Diversity assessment of tree species in Sitio Dicasalarin, Barangay Zabali, Baler, Aurora, Philippines. *Open Journal of Ecology*, *10*(11), 717–728. <https://doi.org/10.4236/oje.2020.1011043>
- Dahms, H. U., & Qian, P. Y. (2004). Life histories of the Harpacticoida (Copepoda, Crustacea): A comparison with meiofauna and macrofauna. *Journal of Natural History*, *38*(14), 1725–1734. <https://doi.org/10.1080/0022293031000156321>
- David, V., Sautour, B., Chardy, P., & Leconte, M. (2005). Long-term changes of the zooplankton variability in a turbid environment: The Gironde estuary (France). *Estuarine, Coastal and Shelf Science*, *64*(2–3), 171–184. <https://doi.org/10.1016/j.ecss.2005.01.014>
- Davies, O. A., Abowei, J. F. N., & Tawari, C. C. (2009). Phytoplankton community of Elechi Creek, Niger Delta, Nigeria—a nutrient-polluted tropical creek. *American Journal of Applied Sciences*, *6*(6), 1143–1152. <https://doi.org/10.3844/ajassp.2009.1143.1152>
- Department of Environment and Natural Resources (DENR). (2016). *Water Quality Guidelines and General Effluent Standards of 2016* (DENR Administrative Order 2016-08). <https://pab.emb.gov.ph/wp-content/uploads/2017/07/DAO-2016-08-WQG-and-GES.pdf>

- Divina, M. A. R., Misanes, V. A. P., Aguinaldo, J. D., & Rafanan, N. S. (2023). FISDA: Smart IoT-based fish farming monitoring system. *The Vector: International Journal of Emerging Science, Technology and Management (IJESTM)*, 32(1).
<https://doi.org/10.69566/ijestm.v32i1.305>
- Durmus, T., Balkis-Ozdelice, N., & Tas, S. (2023). Community structure of tintinnids (Protozoa: Ciliophora) in the eutrophic environment (the Golden Horn Estuary, Turkey). *Journal of Sea Research*, 193, 102382.
<https://doi.org/10.1016/j.seares.2023.102382>
- Echeveste, P., Agustí, S., & Dachs, J. (2011). Cell size dependence of additive versus synergetic effects of UV radiation and PAHs on oceanic phytoplankton. *Environmental Pollution*, 159(5), 1307–1316.
<https://doi.org/10.1016/J.ENVPOL.2011.01.023>
- El-Fattah, I. A. A., Ahmed, M., Aal, M., Elghobashy, H., Fitzsimmons, K., & Diab, A. S. (2008). Zooplankton as live food for fry and fingerlings of Nile tilapia (*Oreochromis niloticus*) and catfish (*Clarias gariepinus*) in concrete ponds.
- Fabry, V. J., Seibel, B. A., Feely, R. A., & Orr, J. C. (2008). Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65(3), 414–432. <https://doi.org/10.1093/ICESJMS/FSN048>
- Florendo, P. E. (2003). Benthic Macroinvertebrate Diversity Index as Water Quality Indicator of Govantes River in Vigan City, Ilocos Sur. *The Vector: International Journal of Emerging Science, Technology and Management (IJESTM)*, 12(1).
<https://doi.org/10.69566/IJESTM.V12I1.162>
- Fondriest Environmental Inc. (2014). *Turbidity, Total Suspended Solids & Water Clarity - Environmental Measurement Systems*. Fondriest Environmental, Inc. .
<https://www.fondriest.com/environmental-measurements/parameters/water-quality/turbidity-total-suspended-solids-water-clarity/>
- Gerhardt, A., Bloor, M., & Mills, C. L. (2011). Gammarus: Important Taxon in Freshwater and Marine Changing Environments. *International Journal of Zoology*, 2011(1), 524276.
<https://doi.org/10.1155/2011/524276>
- Gershwin, L.-A. (2016). *Jellyfish, a natural history*. 224.
- Halfter, S., Cavan, E. L., Swadling, K. M., Eriksen, R. S., & Boyd, P. W. (2020). The Role of Zooplankton in Establishing Carbon Export Regimes in the Southern Ocean – A Comparison of Two Representative Case Studies in the Subantarctic Region. *Frontiers in Marine Science*, 7, 567917.
<https://doi.org/10.3389/FMARS.2020.567917/BIBTEX>
- Hays, G. C., Richardson, A. J., & Robinson, C. (2005). Climate change and marine plankton. *Trends in Ecology & Evolution*, 20(6), 337–344.
<https://doi.org/10.1016/J.TREE.2005.03.004>

- Horak, R. E. A., Ruef, W., Ward, B. B., & Devol, A. H. (2016). Expansion of denitrification and anoxia in the eastern tropical North Pacific from 1972 to 2012. *Geophysical Research Letters*, *43*(10), 5252–5260. <https://doi.org/10.1002/2016GL068871>
- Horsburgh, K. J., & Wilson, C. (2007). Tide-surge interaction and its role in the distribution of surge residuals in the North Sea. *Journal of Geophysical Research: Oceans*, *112*(C8). <https://doi.org/10.1029/2006JC004033>
- Ilac, A. G., Foronda, J. M. S., Ayop, A. N., Europa-Morales, A. L. V., & Ruadap, M. E. V. (2024). Exploring Growth of *Gracilaria* sp. using the Raft Culture Method. *ILMU KELAUTAN: Indonesian Journal of Marine Sciences*, *29*(4), 557–566. <https://doi.org/10.14710/IK.IJMS.29.4.557-566>
- Jain, C. K. (2011). Salinity. *Encyclopedia of Earth Sciences Series, Part 3*, 959–959. https://doi.org/10.1007/978-90-481-2642-2_461
- Kibria, G., Nugegoda, D., Fairclough, R., Lam, P., & Bradly, A. (1997). Zooplankton: its biochemistry and significance in aquaculture. *Naga*, *20*(2), 8–14. <https://ideas.repec.org/a/wfi/wfnaga/33364.html>
- Kim, S. Y., Yang, E. J., Gong, J., & Choi, J. K. (2010). Redescription of *Favella ehrenbergii* (Claparède and Lachmann, 1858) Jörgensen, 1924 (Ciliophora: Choreotrichia), with phylogenetic analyses based on small subunit rRNA gene sequences. *The Journal of Eukaryotic Microbiology*, *57*(6), 460–467. <https://doi.org/10.1111/J.1550-7408.2010.00500.X>
- Lasco, R. (2022). Climate Change and Long-Standing Environmental Problems in the Philippines. *Transactions of the National Academy of Science and Technology*, *42*(2020), 1–7. <https://doi.org/10.57043/TRANSNASTPHL.2020.2311>
- Lauritano, C., Procaccini, G., & Ianora, A. (2012). Gene expression patterns and stress response in marine copepods. *Marine Environmental Research*, *76*, 22–31. <https://doi.org/10.1016/J.MARENRES.2011.09.015>
- Loeblich, A. R., & Tappan, H. (1988). *Foraminiferal Genera and Their Classification*. <https://doi.org/10.1007/978-1-4899-5760-3>
- Mamaril, A. C. (2001). Translocation of the clupeid *Sardinella tawilis* to another lake in the Philippines: A proposal and ecological considerations. *Conservation and Ecological Management of Philippine Lakes in Relation to Fisheries and Aquaculture. Proceedings of the National Seminar-Workshop.*, 133–147. <https://repository.seafdec.org/handle/20.500.12066/5147>
- Martin, J. W., & Davis, G. E. (2001). An Updated Classification Of The Recent Crustacea. *Los Angeles: Natural History Museum of Los Angeles County*, *39*, 129.
- Mendoza, J. A., Verzosa, A. L. R., & Ayop, A. N. (2023). Evaluation of Coral Reef Health Status in the Fish Sanctuary of San Esteban, Ilocos Sur. *The Vector: International Journal of*

The Vector: International Journal of Emerging Science, Technology and Management
Volume 33, Issue 1, January - December 2024

Emerging Science, Technology and Management (IJESTM), 32(1).
<https://doi.org/10.69566/IJESTM.V32I1.302>

- Mwagona, P. C., Chengxue, M., & Hongxian, Y. (2018). Seasonal dynamics of Zooplankton functional groups in relation to environmental variables in Xiquanyan Reservoir, Northeast China. *Annales de Limnologie - International Journal of Limnology*, 54, 33. <https://doi.org/10.1051/LIMN/2018024>
- Naung, N. O. (2018). Community of crustacean larvae in Mon coastal water of Myanmar. *Journal of Aquaculture & Marine Biology, Volume 7*(Issue 5), 275–283. <https://doi.org/10.15406/JAMB.2018.07.00220>
- Ohrel, R. L. Jr., & Register, K. M. (2006). *Volunteer estuary monitoring: A methods manual (2nd Ed)*. https://www.epa.gov/sites/default/files/2015-09/documents/2007_04_09_estuaries_monitoruments_manual.pdf
- Okogwu, Okechukwu. I. (2010). Seasonal variations of species composition and abundance of zooplankton in Ehoma Lake, a floodplain lake in Nigeria. *Revista de Biología Tropical*, 58(1), 171–182. http://www.scielo.sa.cr/scielo.php?script=sci_arttext&pid=S0034-77442010000100013&lng=en&nrm=iso&tlng=en
- Pinti, J., Devries, T., Norin, T., Serra-Pompei, C., Proud, R., Siegel, D. A., Kiørboe, T., Petrik, C. M., Andersen, K. H., Brierley, A. S., & Visser, A. W. (2023). Model estimates of metazoans' contributions to the biological carbon pump. *Biogeosciences*, 20(5), 997–1009. <https://doi.org/10.5194/BG-20-997-2023>
- Ramírez, F. C., & Sabatini, M. E. (2000). The occurrence of Calanidae species in waters off Argentina. *Hydrobiologia*, 439(1), 21–42. <https://doi.org/10.1023/A:1004193401931/METRICS>
- Richardson, A. J. (2008). In hot water: zooplankton and climate change. *ICES Journal of Marine Science*, 65(3), 279–295. <https://doi.org/10.1093/ICESJMS/FSN028>
- Santhanam, R. ., Velayutham, P. ., & Jegatheesan, G. . (1989). *A manual of freshwater ecology : an aspect of fishery management* (International Ed). Daya Publishing House.
- Schram, F. R., & Koenemann, S. (2022a). Copepoda. *The Light and Smith Manual: Intertidal Invertebrates from Central California to Oregon: Fourth Edition, Completely Revised and Expanded*, 446–464. <https://doi.org/10.1093/OSO/9780195365764.003.0011>
- Schram, F. R., & Koenemann, S. (2022b). Cumacea. *The Light and Smith Manual: Intertidal Invertebrates from Central California to Oregon: Fourth Edition, Completely Revised and Expanded*, 495–503. <https://doi.org/10.1093/OSO/9780195365764.003.0031>
- Shaira Elyza R., G., Augusto P., C. P., & Mc Rollyn D., V. (2024). Synthesizing the Ecological and Societal Ramifications of Environmental Stressors on Philippine Marine

- Ecosystems: A Comprehensive Literature Review. *International Journal of Current Science Research and Review*, 07(05). <https://doi.org/10.47191/IJCSRR/V7-I5-13>
- Slotwinski, A., Coman, F., & Richardson, A. J. (2014). Introductory guide to zooplankton identification. In *Integrated Marine Observing System-Brisbane*. Integrated Marine Observing System-Brisbane. https://www.eoas.ubc.ca/~swaterma/473-573/Handouts/IntroductoryZooplanktonFieldGuide_2014.pdf
- Smith, P. R., & Chia, F.-S. (2011). Larval development and metamorphosis of *Sabellaria cementarium* Moore, 1906 (Polychaeta: Sabellariidae). *https://doi.org/10.1139/Z85-156*, 63(5), 1037–1049. <https://doi.org/10.1139/Z85-156>
- Smyth, K., & Elliott, M. (2016). Effects of changing salinity on the ecology of the marine environment. *Stressors in the Marine Environment*, 161–174. <https://doi.org/10.1093/ACPROF:OSO/9780198718826.003.0009>
- Stoecker, D. K., Gallager, S. M., Langdon, C. J., & Davis, L. H. (1995). Particle capture by *Favella* sp. (Ciliata, Tintinnina). *Journal of Plankton Research*, 17(5), 1105–1124. <https://doi.org/10.1093/PLANKT/17.5.1105>
- Tait, R. V. ., & Dipper, Frances. (1998). *Elements of Marine Ecology : an Introductory Course*. 473.
- Thushari, G. G. N., & Senevirathna, J. D. M. (2020). Plastic pollution in the marine environment. *Heliyon*, 6(8), e04709. <https://doi.org/10.1016/J.HELIYON.2020.E04709>
- Turner, J. T. (2004). The Importance of Small Planktonic Copepods and Their Roles in Pelagic Marine Food Webs. *Zoological Studies*.
- United States Environmental Protection Agency. (2006). National Estuary Program coastal condition report. *Library of Congress, Washington, D.C. 20540 USA*. <https://lccn.loc.gov/2007395644>
- Uriarte, I., & Villate, F. (2004). Effects of pollution on zooplankton abundance and distribution in two estuaries of the Basque coast (Bay of Biscay). *Marine Pollution Bulletin*, 49(3), 220–228. <https://doi.org/10.1016/J.MARPOLBUL.2004.02.010>
- Uye, S., Aoto, I., & Onbé, T. (2002). Seasonal population dynamics and production of *Microsetella norvegica*, a widely distributed but little-studied marine planktonic harpacticoid copepod. *Journal of Plankton Research*, 24(2), 143–153. <https://doi.org/10.1093/PLANKT/24.2.143>
- van Velzen, R. (2022). *Salinity - Water Rangers*. Water Rangers. <https://waterrangers.com/2022/04/07/salinity/?v=da984e42a589>
- Villanoy, C. L., Cabrera, O. C., Yñiguez, A., Camoying, M., de Guzman, A., David, L. T., & Lament, P. F. (2011). Monsoon-driven coastal upwelling off Zamboanga peninsula,

The Vector: International Journal of Emerging Science, Technology and Management
Volume 33, Issue 1, January - December 2024

Philippines. *Oceanography*, 24(1), 156–165.

<https://doi.org/10.5670/OCEANOG.2011.12>

Weber-Scannell, P. K., & Duffy, L. K. (2007). Effects of Total Dissolved Solids on Aquatic Organisms: A Review of Literature and Recommendation for Salmonid Species. *American Journal of Environmental Sciences*, 3(1), 1–6.

<https://doi.org/10.3844/AJESSP.2007.1.6>

Yamani, F. Yousif., Skryabin, Valeriy., Gubanova, Aleksandra., Khvorov, Sergey., & Prusova, Irina. (2011a). Marine zooplankton practical guide volume 1. In *Kuwait Institute for Scientific Research*. Kuwait Institute for Scientific Research.

https://www.academia.edu/43277931/MARINE_ZOOPLANKTON_PRACTICAL_GUIDE_Volume_1

Yamani, F. Yousif., Skryabin, Valeriy., Gubanova, Aleksandra., Khvorov, Sergey., & Prusova, Irina. (2011b). Marine zooplankton practical guide volume 2. In *Kuwait Institute for Scientific Research*. Kuwait Institute for Scientific Research.

https://www.academia.edu/43277726/MARINE_ZOOPLANKTON_PRACTICAL_GUIDE_Volume_2

Zheng, Z., Fu, Y., Liu, K., Xiao, R., Wang, X., & Shi, H. (2018). Three-stage Vertical Distribution of Seawater Conductivity. *Scientific Reports 2018 8:1*, 8(1), 1–10.

<https://doi.org/10.1038/s41598-018-27931-y>